



Green Bay • Fox Valley • Marinette

2223 Lime Kiln Road, Green Bay, WI 54311 • Phone: 920-430-8113 • Fax: 920-430-8124

osmsgb.com

Physical Therapy Protocol: Distal Biceps Repair

Philosophy:

Distal biceps rupture is an injury most commonly seen in men, usually during middle age, although it can truly happen to anyone. The most common mechanism of injury is a forced elbow extension while the muscle is tensioned in flexion. A distinct “pop” is often felt at the elbow, and a soft retraction of the tendon is palpable into the upper arm area. Loss of supination strength and even elbow flexion strength is common with unrecognized injuries. If your doctor does recommend surgery, here is a guideline for recommended treatment following the repair procedure. Initially, immobilization will be held at flexion, followed by gradual return to a neutral position. A strong, pain-free elbow with functional range-of-motion only comes about through an anatomic reconstruction of the biceps with aggressive (but appropriate) physical therapy performed with the therapist and at home.

Phase I, surgery to 6 weeks

OSMS appointments:

Medical appointments at 1 and 6 weeks with films at first visit

Physical therapy appointments begin at first visit, and continue every 5-7 days

Rehabilitation Goals:

Achieve excellent hand and shoulder range-of-motion

Reduce swelling in the limb

Control pain

Precautions:

Posterior elbow splint at 90 degrees with forearm neutral

Range-of-Motion Exercises:

Post-surgical splint for 7 days

After seven days, elbow is placed in hinged IROM with brace unlocked at 45° to full flexion.

Week 2: 45° to full flexion

Week 4: 30° to full flexion

Week 5: 20° to full flexion

Week 6: 10° to full flexion

Week 8: Unlock completely and discontinue brace if adequate motor control is achieved

Suggested Therapeutic Exercises:

Ball squeezes, submaximal shoulder isometrics with forearm in neutral

Cardiovascular Exercises:

Lower body circuit training or lower body cardio



Green Bay • Fox Valley • Marinette

2223 Lime Kiln Road, Green Bay, WI 54311 • Phone: 920-430-8113 • Fax: 920-430-8124

osmsgb.com

Progression Criteria:

Patient may progress to phase II after 4 weeks if they have pain well-controlled

Phase II, (after Phase I criteria met, usually at 6-8 weeks)

OSMS appointments:

Medical appointments at 6 weeks

Physical therapy appointments continue at every 5-7 days or so

Rehabilitation Goals:

Restore full ROM and adequate motor control

Precautions:

No biceps strength training outside of supervised PT visits

Range-of-Motion Exercises:

Continue progression as outlined

Week 6: 10 to full flexion

Week 8: Unlock completely and discontinue brace if adequate motor control is achieved

Suggested Therapeutic Exercises:

Begin muscle strengthening exercises for wrist and forearm

Shoulder strengthening

Progression Criteria:

Patient may progress to phase III after 2 weeks if they have met the above stated goals & have pain well-controlled.

Phase III, (after Phase II criteria met, usually 8-16 weeks)

OSMS appointments:

Medical appointment at 12 weeks

Physical therapy appointments adjust to every 10-14 days, advance to home program

Rehabilitation Goals:

PREs for elbow flexion, extension, supination, and pronation

Allow light biceps training from 12-14 weeks

Initiate endurance program to simulate desired work or sport requirements

Begin a return to sport specific exercises

Full resistance biceps curls allowed at 16 weeks



Green Bay • Fox Valley • Marinette

2223 Lime Kiln Road, Green Bay, WI 54311 • Phone: 920-430-8113 • Fax: 920-430-8124

osmsgb.com

References:

- Greenberg, J.A., Fernandez J.J., Wang T, and Turner C, (2003) Endobutton-assisted repair of distal biceps tendon ruptures. *J. Shoulder Elbow Surgery* 12: 484.
- Hand and Upper Limb Clinic Protocol ;(2010)
http://www.hulc.ca/pdf/Distal_Bicep_Protocol_2010.pdf .
- Indiana Hand Diagnosis and Treatment Manual for Physicians and Therapists, 4th Edition, (2001).
- Rose, D.M., Archibald, J.D., Sutter, E.G. et al. (2011) Biomechanical analysis suggests early rehabilitation is possible after single-incision EndoButton distal biceps repair with FiberWire. *Knee Surg Sports Traumatol Arthrosc* 19: 1019. doi:10.1007/s00167-010-1348-1.
- [Smith, J.R.](#), Amirfeyz, R. (2016) Does immediate elbow mobilization after distal biceps tendon repair carry the risk of wound breakdown, failure of repair, or patient dissatisfaction? *J Shoulder Elbow Surg.* 2016 May;25(5):810-5. doi: 10.1016/j.jse.2015.11.066.
- Spencer E., Tisdale A., Kostka K., and Ivy R. (2008) Is therapy necessary after distal biceps tendon repair? *Hand (N.Y.) American Association for Hand Surgery* 2008 Dec, 3(4): 316-319
- UW Health Hand and Upper Extremity Clinic Protocol.
- *Clinical Orthopaedic Rehabilitation: A team approach*, 4th edition, (2018) 26, 164-173. E1.