

Physical Therapy Protocol: Biceps Tenodesis

Philosophy:

Our biceps tenodesis protocol begins with early flexibility and stretching, followed by a progression to strengthening, and then advanced function. These are progressed quickly to prevent the most common cause of failure (stiffness). Individual variations will occur depending on surgeon input, concomitant rotator cuff findings, and patient response to treatment. The following are guidelines for biceps tenodesis rehabilitation.

Phase I, surgery to 4 weeks

OSMS appointments:

- Medical appointment at 2 weeks with films
- Rehabilitation will begin as directed by your physician and as indicated on your rehabilitation order

Rehabilitation Goals:

- Cryotherapy unit to the shoulder: twenty minutes every two hours to reduce swelling
- Full passive ROM for shoulder flexion, abduction, IR and ER
- Achieve activation of the stabilizing muscles for the GH and scapulothoracic joints

Precautions:

- Strict sling for 4 weeks
- No lifting anything heavier than 12 oz
- No bicep tension for 4 weeks to protect tenodesis. This includes avoiding long arm flexion and resisted supination or elbow flexion (No ER past 40°, no shoulder extension).

Suggested Therapeutic Exercises:

- Grip strengthening
- Forearm and wrist ROM
- Cervical spine & scapular active ROM
- Desensitization for axillary n distribution
- Elbow AAROM/PROM without resistance (to allow the biceps tendon time to heal into the new insertion site on the humerus without stress)
- AAROM/PROM for shoulder flexion, abduction, IR and ER

Cardiovascular Exercises (with sling on):

- Walking, Stairmaster, or stationary bike



Progression Criteria:

- Incisional healing, full PROM of shoulder, and of elbow in sup/pron/flex/extension
- Negative impingement pain or shoulder apprehension

Phase II, 4-6 weeks

OSMS appointments:

- MD appointment at 6 weeks
- Rehabilitation appointments every 5-7 days

Rehabilitation Goals:

- Advance to AROM for shoulder flexion, abduction, IR and ER to neutral (and normal scapulothoracic movement)
- Begin proprioceptive and dynamic neuromuscular control training
- Strengthen shoulder and scap stabilizers
- Full discontinuation of sling

Precautions:

- No cross-frictional massage
- No swimming, throwing or overhead serves
- Avoid activities that have risk for falls

Suggested Therapeutic Exercises:

- Gentle shoulder mobilizations as needed
- AAROM/AROM in all planes-assessing scapular rhythm
- RTC strengthening
- Ball squeezes
- Core strengthening

Cardiovascular Exercises:

- Stationary bike without arms
- Impact exercises are only allowed once the patient exhibits full RTC strengthening in neutral position (to avoid distractive forces when landing)

Progression Criteria:

- Full shoulder AROM, scapular posture at rest and dynamic scapular control with ROM and/or functional activities

Phase III, 6-12 weeks

OSMS appointments:

- MD appointment at 12 weeks
- Rehabilitation appointments every 1-2 weeks

Rehabilitation Goals:

- Full shoulder and scapular ROM, all planes
- 5/5 RTC strength and peri-scapular strength

Precautions:

- No swimming, throwing, or overhead serves until 12 weeks
- Avoid activities that have risk for falls

Suggested Therapeutic Exercises:

- At 6 weeks, begin light isometrics with the arm at side
- At 6 weeks, begin scapular PREs
- At 8 weeks, UE ergometer, weighted ball toss, body blade, for eccentrically resisted motion
- Demonstrate stability with higher velocity movements and change of direction that replicate sport-specific patterns
- Pain-free return to high velocity overhead movements
- Improve core and hip strength & flexibility to eliminate any compensatory stresses to the shoulder
- Work capacity endurance for specific demands

Progression Criteria:

- The patient can progress to full activities when they have met goals pain-free (appropriate RTC and scapular musculature performance for chest level activities)
- Return to throwing and begin swimming at 12 weeks
- Throw from pitcher's mound at 18 weeks
- Return to collision sports at 26 weeks
- MMI is usually 26 weeks



Green Bay • Fox Valley • Marinette • Oshkosh

Phone: 920-430-8113 • Fax: 920-593-2986

osmsgb.com

References:

- Cools AM, Dewitte V, Lanszweert F, Notebaert D, Roets A, Soetens B, Cagnie B, Witvrouw EE. Rehabilitation of Scapular Muscle Balance: Which Exercises to Prescribe?. *Am. J. Sports Med.* July 2007; 35(10); 1744-1751.
- Krupp RJ, Kevern MA, Gaines MD, Kotara S, Singleton SB. Long Head of the Biceps Tendon Pain: Differential Diagnosis and Treatment. *Jour Ortho & Sports PT.* Feb 2009; 39(2): 55-70.
- Nho SJ, Strauss EJ, Lenart BA, Provencher MT, Mazzocca AD, Verma NN, Romeo AA. Long head of the biceps tendiopathy: diagnosis and management. *J Am Acad Orthop Surg.* 2010; 18(10): 645-656.
- Reinold MM, Escamilla R, Wilk KE. Current concepts in the scientific and clinical rationale behind exercises for glenohumeral and scapulothoracic musculature. *J Orthop and Sports Medicine.* 2009; 39(2): 105-117.
- Romeo AA, Mazzocca AD, Tauro JC. Arthroscopic Biceps Tenodesis. *Arthroscopy.* Feb 2004; 20(2): 206-213.
- Ryu JH, Pedowitz RA. Rehabilitation of biceps tendon disorders in athletes. *Clinics in Sports Medicine.* 2010; 29: 229-46.
- Wilk KE, Meister K, Andrews JR. Current concepts in the rehabilitation of the overhead throwing athlete. *Am J Sports Med.* 2002;30(1):136-151.
- Wilk KE, Yenchak AJ, Arrigo CA, Andrews JR. The advanced throwers ten exercise program: a new exercise series for enhanced dynamic shoulder control in the overhead throwing athlete. *Phys Sportsmed.* 2011;39(4):90-97