



Green Bay • Fox Valley • Marinette

2223 Lime Kiln Road, Green Bay, WI 54311 • Phone: 920-430-8113 • Fax: 920-430-8124

osmsgb.com

Iliopsoas Lengthening after Total Hip Arthroplasty Physical Therapy Protocol

Please use appropriate clinical judgment during all exercise progressions. The specific exercises given in this protocol are provided for guidance, but it is important to use clinical judgment when determining appropriate progressions with the physician provided WBing and ROM restrictions. Any questions/concerns, please do not hesitate to contact the office at (920)430-8113.

Weightbearing: The patient is WBAT with assistive device for 2 weeks post-operatively, then will be cleared to wean off assistive device as tolerated.

PHASE I, Immediate Rehabilitation (weeks 1-6)

Begin therapy post-operative day #1-7

Goals:

- Protection of lengthened tissue
- Restore ROM within guidelines
- Prevent muscular inhibition and gait abnormalities
- Diminish pain and inflammation

Precautions:

- Do NOT aggressively push through pain/pinching
- Progress cautiously with signs of hip flexor tendonitis, trochanteric bursitis, and synovitis

PROM Guidelines:

- All planes as tolerated; do not force range through pain or pinching. Initiate iliopsoas stretching (gentle and pain-free) as soon as patient can tolerate.

Weeks 1-4:

- PROM to hip within ROM guidelines- avoid pain/pinch. Do long axis traction with circumduction (for circulating synovial fluid and, therefore, cartilage health)
- Add stationary bike without resistance at 2-3 weeks, maximizing seat height to avoid psoas irritation
- Exercises: hamstring stretch, single knee to chest, prone quad stretch, supine hip flexor stretch, quad sets, ankle pumps, glut sets, transverse abdominal activation, hip adduction/abduction isometrics



Green Bay • Fox Valley • Marinette

2223 Lime Kiln Road, Green Bay, WI 54311 • Phone: 920-430-8113 • Fax: 920-430-8124

osmsgb.com

- Manual Considerations: scar massage, STM/MFR to: TFL, ITB, psoas, iliacus, hip adductors, quadratus lumborus, paraspinals
- Modalities for pain control and swelling

Weeks 4-6:

- Continue with previous exercises and manual
- PROM: Continue to restore full ROM as tolerated
- Exercises: sub-max, pain-free hip flexion- avoid hip flexor tendonitis; quad and hamstring isotonic exercise. Progress CKC exercises as tolerated, focusing on core and hip stability and maintaining good hip/knee/ankle alignment
- Manual Considerations: scar massage, STM/MFR to: TFL, ITB, psoas, iliacus, hip adductors, quadratus lumborum, paraspinals
- Modalities for pain control and swelling

PHASE II, Intermediate Rehabilitation (weeks 6-12)

Weeks 6-8:

- Continue with previous exercises
- PROM: Continue to restore full ROM as tolerated
- Exercises: Progress core strengthening, CKC as tolerated, may advance walking program, stationary bike with resistance, elliptical
 - Progress hip PREs as tolerated: leg press, hamstring curls, leg extensions, squat progression
 - Hip hiking, eccentric step downs
 - Step ups: forward and side as tolerated; progress to diagonal
- Manual Considerations: As needed

PHASE III, Advanced Rehabilitation

Weeks 12 and beyond:

- Continue with previous exercises and manual
- PROM: Continue to restore full ROM as tolerated
- Exercises: Progress all strengthening as tolerated. Use clinical judgement to determine lunge/jump/hop/jog training appropriate for patient lifestyle and goals



Green Bay • Fox Valley • Marinette

2223 Lime Kiln Road, Green Bay, WI 54311 • Phone: 920-430-8113 • Fax: 920-430-8124

osmsgb.com